





2. Raphael M, Jaeger M and van Vlymen J. Easily Adoptable Total Joint Arthroplasty Program Allows Discharge Home in Two Days. *Can J Anesth.* 2011; 58: 902-910. [\[Crossref\]](#)
3. Lombardi AV, Viacava AJ and Berend KR. Rapid Recovery Protocols and Minimally Invasive Surgery Help Achieve High Knee Flexion. *Clin Orthop Rel Res.* 2006; 452: 117-122. [\[Crossref\]](#)
4. Berger RA, Jacobs JJ, Meneghini RM, Della Valle C, Paprosky W, Rosenberg AG, et al. Rapid Rehabilitation and Recovery with Minimally Invasive Total Hip Arthroplasty. *Clin Orthop Rel Res.* 2004; 429: 239-247. [\[Crossref\]](#)
5. Williams-Russo P, Sharrock NE, Mattis S, Liguori GA, Mancuso C, Peterson MG, et al. Randomized trial of hypotensive epidural anesthesia in older adults. *Anesthesiology.* 1999; 91:926-935. [\[Crossref\]](#)
6. Cram P, Cai X, Lu X, Vaughan-Sarrazin MS and Miller BJ. Total Knee Arthroplasty Outcomes in Top-Ranked and Non-Top-Ranked Orthopedic Hospitals: An Analysis of Medicare Administrative Data. *Mayo Clin Proc.* 2012 ; 87: 341-348. [\[Crossref\]](#)
7. Kirksey M, Chiu YL, Ma Y, Della Valle AG, Poulosides L, Gerner P, et al. Trends in In-Hospital Major Morbidity and Mortality After Total Joint Arthroplasty: United States 1998-2008. *Anesth Analg.* 2012 ; 115: 321-327. [\[Crossref\]](#)
8. Rodgers A, Walker N, Schug S, McKee A, Kehlet H, van Zundert A, et al. Reduction of Postoperative Mortality and Morbidity with Epidural or Spinal Anesthesia: Results from Overview of Randomised Trials. *BMJ.* 2000; 321: 1493. [\[Crossref\]](#)
9. Mauerman WJ, Shilling AM and Zuo Z. A comparison of neuraxial block versus general anesthesia for elective total hip replacement: a meta-analysis. *Anesth Analg.* 2006; 103:1018-1025. [\[Crossref\]](#)
10. Macfarlane AJ, Prasad GA, Chan VW and Brull R. Does regional anaesthesia improve outcome after total hip arthroplasty? A systematic review. *Br J Anaesth.* 2009; 103: 335-345. [\[Crossref\]](#)
11. Gonano C, Leitgeb U, Sitzwohl C, Ihra G, Weinstabl C, Kettner SC, et al. Spinal Versus General Anesthesia for Orthopedic Surgery: Anesthesia Drug and Supply Costs. *Anesth Analg.* 2006; 102: 524-529. [\[Crossref\]](#)
12. Brull R, McCartney CJ, Chan VW and El-Beheiry H. Neurological complications after regional anesthesia: contemporary estimates of risk. *Anesth Analg.* 2007; 104: 965-974. [\[Crossref\]](#)
13. Cushner FD. Do Regional Analgesia and Peripheral Blocks Still Have a Place in Joint Arthroplasty? *Am J Orthopedics.* 2015; 44: 9-12. [\[Crossref\]](#)
14. Widmer B, Lustig S, Scholes CJ, Molloy A, Leo SP, Coolican MR, et al. Incidence and severity of complications due to femoral nerve blocks performed for knee surgery. *Knee.* 2013; 20: 181-185. [\[Crossref\]](#)
15. Pelt CE, Anderson AW, Anderson MB, Van Dine C and Peters CL . Postoperative falls after total knee arthroplasty in patients with a femoral nerve catheter: can we reduce the incidence? *J Arthroplasty.* 2014; 29: 1154-1157. [\[Crossref\]](#)
16. Sharma S, Richard Iorio, Lawrence M. Specht, Sara Davies-Lepie and William L. Healy . Complications of femoral nerve block for total knee arthroplasty. *Clin Orthop Relat Res.* 2010 ; 468: 135-140. [\[Crossref\]](#)
17. Toftdahl K, Nikolajsen L, Haraldsted V, Madsen F, Tønnesen EK, Søballe K, et al. Comparison of peri- and intraarticular analgesia with femoral nerve block after total knee arthroplasty: a randomized clinical trial. *Acta Orthop.* 2007; 78:172-179. [\[Crossref\]](#)
18. Busch CA, Shore BJ, Bhandari R, Ganapathy S, MacDonald SJ, Bourne RB, et al. Efficacy of periarticular multimodal drug injection in total knee arthroplasty. A randomized trial. *J Bone Joint Surg Am.* 2006; 88:959-963. [\[Crossref\]](#)
19. Teng Y, Jiang J, Chen S, Zhao L, Cui Z, Khan MS, et al . Periarticular multimodal drug injection in total knee arthroplasty. *Knee Surg Sports Traumatol Arthrosc.* 2014; 22:1949-1957. [\[Crossref\]](#)
20. Calum HC. Arthur, Alexander MW, Charles Leeson-Payne and Steffen J. Breusch. Comparison of Peri-Articular Multimodal Drug Injection with Femoral and Sciatic Nerve Block after Total Knee Arthroplasty. *Open J Ortho.* 2012; 2:155-158. [\[Crossref\]](#)
21. Parvataneni HK, Shah VP, Howard H, Cole N, Ranawat AS, Ranawat CS, et al. Controlling Pain After Total Hip and Knee Arthroplasty Using a Multimodal Protocol with Local Periarticular Injections: A Prospective Randomized Study. *J Arthroplasty.* 2007; 22: 33-38. [\[Crossref\]](#)
22. Mark J. Spangehl, Henry D. Clarke, Joseph G. Hentz, Lopa Misra, Joshua L. Blocher, David P. Seamans, et al. The Chitranjan Ranawat Award: Periarticular Injections and Femoral and Sciatic Blocks Provide Similar Pain Relief after TKA: a Randomized Clinical Trial. *Clin Orthop Relat Res.* 2015; 473: 45-53. [\[Crossref\]](#)
23. Emerson RH. Comparison of Infiltration with Long-Acting Bupivacaine to a Femoral Nerve Catheter for Total Knee Replacement. American Academy of Orthopaedic Surgeons Annual Meeting; March 2014; New Orleans, LA. Poster P124. [\[Crossref\]](#)
24. Affas F, Nygård EB, Stiller CO, Wretenberg P and Olofsson C. Pain control after total knee arthroplasty: a randomized trial comparing local infiltration anesthesia and continuous femoral block. *Acta Orthop.* 2011; 82:441-447. [\[Crossref\]](#)
25. Chaumeron A, Audy D, Drolet P, Lavigne M and Vendittoli PA. Periarticular injection in knee arthroplasty improves quadriceps function. *Clin Orthop Relat Res.* 2013; 471:2284-2295. [\[Crossref\]](#)
26. Ng FY, Ng JK, Chiu KY, Yan CH and Chan CW. Multimodal periarticular injection vs continuous femoral nerve block after total knee arthroplasty: a prospective, crossover, randomized clinical trial. *J Arthroplasty.* 2012;27:1234-1238. [\[Crossref\]](#)
27. Widmer BJ, Scholes CJ, Pattullo GG, Oussedik SI, Parker DA, Coolican MR, et al. Is femoral nerve block necessary during total knee arthroplasty?: a randomized controlled trial. *J Arthroplasty.* 2012; 27:1800-1805. [\[Crossref\]](#)
28. Pulido L, Ghanem E, Joshi A, Purtill JJ and Parvizi J. Periprosthetic Joint Infection: The Incidence, Timing, and Predisposing Factors. *Clin Orthop Rel Res.* 2008; 466: 1710-1715. [\[Crossref\]](#)
29. Bierbaum BE, Callaghan JJ, Galante JO, Rubash HE, Tooms RE, Welch RB, et al. An Analysis of Blood Management in Patients Having a Total Hip or Knee Arthroplasty. *J Bone Joint Surg Am.* 1999; 81: 2 -10. [\[Crossref\]](#)
30. Nelson CL and Bowen WS. Total hip arthroplasty in Jehovah's Witnesses without blood transfusion. *J Bone Joint Surg Am.* 1986; 68: 350 -353. [\[Crossref\]](#)
31. Moonen AF, Neal TD and Pilot P. Peri-operative blood management in elective orthopaedic surgery. A critical review of the literature. *Injury.* 2006; 37: 11-16. [\[Crossref\]](#)
32. Jeulsgaard P, Larsen UT, Sørensen JV, Madsen F and Søballe K. Hypotensive Epidural Anesthesia in Total Knee Replacement Without Tourniquet: Reduced Blood Loss and Transfusion. *Reg Anesth Pain Med.* 2001; 26: 105-110. [\[Crossref\]](#)
33. Hiippala ST, Trid LJ, Wennerstrand MI, Arvela JV, Niemelä HM, Mäntylä SK, et al. Tranexamic Acid Radically Decreases Blood Loss and Transfusions Associated with Total Knee Arthroplasty. *Anesth Analg.* 1997; 84: 839-844. [\[Crossref\]](#)
34. Ekback G, S Axelsson K, Rytberg L, Edlund B, Kjellberg J, Weckström J, et al. Tranexamic Acid Reduces Blood Loss in Total Hip Replacement Surgery. *Anesth Analg.* 2000; 91: 1124-1130. [\[Crossref\]](#)
35. Husted H, Bønd L, Sonne-Holm S, Holm G, Jacobsen TW, Gebuhr P, et al. Tranexamic acid reduces blood loss and blood transfusions in primary total hip arthroplasty: A prospective randomized double-blind study in 40 patients. *Acta Orthop Scand.* 2003; 74: 665-669. [\[Crossref\]](#)
36. Niskanen RO and Korkala O. Od loss in cemented hip arthroplasty: A randomized, double-blind study of 39 patients. Tranexamic acid reduces blood loss with osteoarthritis. *Acta Orthop.* 2005; 76: 829-832. [\[Crossref\]](#)
37. Ho KM and Ismail H. Use of intravenous tranexamic acid to reduce allogeneic blood transfusion in total hip and knee arthroplasty: A meta-analysis. *Anesth Intensive Care.* 2003; 31: 529-37. [\[Crossref\]](#)
38. Hill C, Flamant R, Mazas F and Evrard J. Prophylactic cefazolin versus placebo in total hip replacement. Report of a multicentre double-blind randomised trial. *Lancet.* 1981; 1:795-796. [\[Crossref\]](#)
39. Dellinger EP, Gross PA, Barrett TL, Krause PJ, Martone WJ, McGowan JE Jr, et al. Quality standard for antimicrobial prophylaxis in surgical procedures: The Infectious Diseases Society of America. *Infect Control Hosp Epidemiol.* 1994;15:182-188. [\[Crossref\]](#)
40. Ritter MA. Operating Room Environment. *Clin Orthop Relat Res.* 1999:103-109. [\[Crossref\]](#)
41. Sossai D, D. Sossai, G. Dagnino, F. Sanguineti, and F. Franchin. Mobile laminar air flow screen for additional operating room ventilation: reduction of intraoperative bacterial contamination during total knee arthroplasty. *J Orthop Traumatol.* 2011; 12: 207-211. [\[Crossref\]](#)
42. Smith EB, Raphael JJ, Maltenfort MG, Honsawek S, Dolan K, Younkins EA, et al . The effect of laminar air flow and door openings on operating room contamination. *J Arthroplasty.* 2013; 28:1482-1485. [\[Crossref\]](#)
43. Pokrywka M and Byers K. Traffic in the operating room: a review of factors influencing air flow and surgical wound contamination. *Infect Disord Drug Targets.* 2013; 13:156-161. [\[Crossref\]](#)
44. Panahi P, Stroh M, Casper DS, Parvizi J and Austin MS. Operating room traffic is a major concern during total joint arthroplasty. *Clin Orthop Relat Res.* 2012 ;470: 2690-2694. [\[Crossref\]](#)
45. Bédard M, Pelletier-Roy R, Angers-Goulet M, Leblanc PA and Pelet S. Traffic in the operating room during joint replacement is a multidisciplinary problem. *Can J Surg.* 2015; 58:232-236. [\[Crossref\]](#)